

## Coastal wet tropospheric correction: GPD vs RWT

Study variable	<b>GPD</b>
Reference variable	<b>RWT</b>
Missions	Envisat ( <i>en</i> )
Period	[19259, 22209]

Creation date : 2011/08/30

### Contents

<b>A001</b>	<b>4</b>
<b>A002</b>	<b>5</b>
<b>A003</b>	<b>6</b>
<b>A004</b>	<b>8</b>
<b>A101</b>	<b>9</b>
<b>A102</b>	<b>10</b>
<b>A103</b>	<b>11</b>
<b>A104</b>	<b>12</b>
<b>A201</b>	<b>13</b>
<b>A202</b>	<b>18</b>
<b>A203</b>	<b>20</b>
<b>A204</b>	<b>23</b>
<b>A205</b>	<b>25</b>
<b>A206</b>	<b>27</b>
<b>A207</b>	<b>30</b>

<b>A208</b>	<b>31</b>
<b>A209</b>	<b>32</b>
<b>C001</b>	<b>33</b>
<b>C002</b>	<b>34</b>
<b>C003</b>	<b>35</b>
<b>C004</b>	<b>36</b>

## Study overview

In this study, the wet tropospheric correction computed by the University of Porto, Faculty of science in the scope of the Sea level CCI project (WP2710) has been compared with the radiometric correction to calculate the Envisat sea-level height (SSH).

The impact of using these wet tropospheric corrections on the SSH computation has been analyzed for Envisat mission from September 2002 (cycle 10) to October 2010 (Cycle 93).

The major aim of WP2710 is to provide a wet tropospheric correction for the coastal zone, applicable to all missions, fully compatible with respect to the microwave radiometer (MWR) based correction that shall be adopted in the open ocean, and ensuring its continuity and consistency in the open ocean/coastal transition zone. It has been produced by the university of Porto, Faculty of science (J. Fernandes).

This study has been performed on points where the studied correction is a valid estimate (GPD flag=1) and on non corrupted ocean points where it equals the radiometric correction (GPD flag=0).

For Envisat mission, the reference correction is the radiometric wet tropospheric correction present in GDR products.

All the validation diagnostics displayed in this report have been performed in agreement with the Sea-Level CCI Product Validation Plan (PVP).



## Diagnostic A001 (mission en)

**Name :** Temporal evolution of differences between both altimetric components

**Input data :** Along-track altimetric components

**Description :** The temporal evolution of global statistics (mean, variance, slope) of differences between 2 different standards of a same altimetric component (sea surface height correction, altimeter parameter, orbit) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) . These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Global internal analyses



## Diagnostic A002 (mission en)


**Name :** Map of differences between both altimetric components over all the period

**Input data :** Along-track altimetric components


**Description :** The map of global statistics (mean, standard deviation) of differences between 2 different standards of a same altimetric component (sea surface height correction, altimeter parameter, orbit) are calculated over a given period which is the longer as possible to have obtain reliable statically results. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Global internal analyses

Mean of GPD – RWT  
Mission en, cycles 9 to 93



Mean ( cm )  
Standard deviation of GPD – RWT  
Mission en, cycles 9 to 93





## Diagnostic A003\_a (mission en)

**Name :** Periodogram derived from temporal evolution of altimetric component differences

**Input data :** Along-track altimetric components

**Description :** The periodogram derived from temporal and global altimetric component differences is calculated from cycle by cycle monitoring of altimetric component differences (derived from diagnostic A001). It is calculated from the mean or the variance differences. The Periodogram can be calculated for all the periods, but it can be focused on a dedicated period.

Diagnostic type : Global internal analyses





## Diagnostic A003\_b (mission en)

**Name :** Periodogram derived from temporal evolution of altimetric component differences

**Input data :** Along-track altimetric components

**Description :** The periodogram derived from temporal and global altimetric component differences is calculated from cycle by cycle monitoring of altimetric component differences (derived from diagnostic A001). It is calculated from the mean or the variance differences. The Periodogram can be calculated for all the periods, but it can be focused on a dedicated period.

Diagnostic type : Global internal analyses




### Diagnostic A004 (mission en)

**Name :** Altimetric component differences versus coastal distances

**Input data :** Along-track altimetric components

**Description :** Mean and standard deviation of the differences between 2 different standards of a same altimetric component (sea surface height correction, altimeter parameter, orbit) are computed and plotted in function of coastal distances between 0 and 100 km.

Diagnostic type : Global internal analyses





## Diagnostic A101 (mission en)

**Name :** Temporal evolution of SSH crossovers

**Input data :** Sea Surface Height (SSH) crossovers

**Description :** The temporal evolution of global statistics (mean, standard deviation) of SSH differences are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Global internal analyses




## Diagnostic A102 (mission en)

**Name :** Differences between temporal evolution of SSH crossovers

**Input data :** Sea Surface Height (SSH) crossovers

**Description :** The difference of temporal evolution between the global statistics (mean, standard deviation) of SSH differences are calculated using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Global internal analyses



### Diagnostic A103 (mission en)


**Name :** Map of SSH crossovers

**Input data :** Sea Surface Height (SSH) crossovers


**Description :** The differences between maps of SSH crossovers differences (mean, variance) are calculated using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Global internal analyses

Mean of SSH with GPD  
Mission en, cycles 9 to 93



Mean ( cm )  
Mean of SSH with RWT  
Mission en, cycles 9 to 93




**Diagnostic A104 (mission en)**

**Name :** Differences between maps of SSH crossovers

**Input data :** Sea Surface Height (SSH) crossovers

**Description :** The differences between maps of SSH crossovers (derived from diagnostic A103) are calculated from the SSH crossover differences (mean, standard deviation) using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

**VAR(SSH with GPD) – VAR(SSH with RWT)**  
**Mission en, cycles 9 to 93**



-2      -0.8      0.4      1.6

**SSH crossovers : difference of variances ( cm<sup>2</sup> )**


### Diagnostic A201 a (mission en)

**Name :** Temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) , or separating North and South hemispheres.

Diagnostic type : Global internal analyses





## Diagnostic A201\_b (mission en)

**Name :** Temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) , or separating North and South hemispheres.

Diagnostic type : Global internal analyses




## Diagnostic A201\_c (mission en)

**Name :** Temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) , or separating North and South hemispheres.

Diagnostic type : Global internal analyses



## Diagnostic A201\_d (mission en)


**Name :** Temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) , or separating North and South hemispheres.

Diagnostic type : Global internal analyses

Global MSL  
Mission en, cycles 9 to 93



## Diagnostic A201\_e (mission en)


**Name :** Temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA


**Description :** The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) , or separating North and South hemispheres.

Diagnostic type : Global internal analyses

Global MSL, selecting even pass numbers  
Mission en, cycles 9 to 93



Global MSL, selecting odd pass numbers  
Mission en, cycles 9 to 93




### Diagnostic A202.a (mission en)

**Name :** Differences between temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The differences between temporal evolution of SLA are calculated from statistics derived from diagnostic A201 (mean, variance) using 2 different components in the SLA calculation. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) or separating North and South hemispheres.

Diagnostic type : Global internal analyses





## Diagnostic A202\_b (mission en)

**Name :** Differences between temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The differences between temporal evolution of SLA are calculated from statistics derived from diagnostic A201 (mean, variance) using 2 different components in the SLA calculation. They are calculated globally, but also separating ascending and descending passes (except for SLA Grids) or separating North and South hemispheres.

Diagnostic type : Global internal analyses



## Diagnostic A203\_a (mission en)


**Name :** Map of Sea Level Anomaly (SLA) over all the period

**Input data :** Along track SLA

**Description :** The map of global statistics (mean, standard deviation) of SLA are calculated using successively both altimetric components in the SLA calculation over a large period. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.


Diagnostic type : Global internal analyses

SLA with GPD trends  
Mission en, cycles 9 to 93



Trends (mm/yr)

SLA with RWT trends  
Mission en, cycles 9 to 93



Trends (mm/yr)

## Diagnostic A203\_b (mission en)


**Name :** Map of Sea Level Anomaly (SLA) over all the period

**Input data :** Along track SLA

**Description :** The map of global statistics (mean, standard deviation) of SLA are calculated using successively both altimetric components in the SLA calculation over a large period. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.


Diagnostic type : Global internal analyses

SLA with GPD trends : even pass numbers  
Mission en, cycles 9 to 93



-20.26998 -8.22361 3.82277 15.86915  
Trends (mm/yr)

SLA with RWT trends : even pass numbers  
Mission en, cycles 9 to 93



-20.12909 -8.12092 3.88726 15.89543  
Trends (mm/yr)

## Diagnostic A203\_c (mission en)


**Name :** Map of Sea Level Anomaly (SLA) over all the period

**Input data :** Along track SLA


**Description :** The map of global statistics (mean, standard deviation) of SLA are calculated using successively both altimetric components in the SLA calculation over a large period. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Global internal analyses

SLA with GPD trends : odd pass numbers  
Mission en, cycles 9 to 93



SLA with RWT trends : odd pass numbers  
Mission en, cycles 9 to 93



**Diagnostic A204 a (mission en)**


**Name :** Differences between maps of SLA

**Input data :** Along track SLA

**Description :** The difference of SLA maps (mean, standard deviation, slope) is calculated from maps derived from diagnostic A203 using successively both altimetric components in the SLA calculation over a given period. This can be done globally, or separating in ascending and descending passes (except for SLA Grids).

Diagnostic type : Global internal analyses

**SLA with GPD trends – SLA with RWT trends**  
**Mission en, cycles 9 to 93**



## Diagnostic A204\_b (mission en)


**Name :** Differences between maps of SLA

**Input data :** Along track SLA


**Description :** The difference of SLA maps (mean, standard deviation, slope) is calculated from maps derived from diagnostic A203 using successively both altimetric components in the SLA calculation over a given period. This can be done globally, or separating in ascending and descending passes (except for SLA Grids).

Diagnostic type : Global internal analyses

SLA with GPD trends – SLA with RWT trends : even pass numbers  
Mission en, cycles 9 to 93



SLA with GPD trends – SLA with RWT trends : odd pass numbers  
Mission en, cycles 9 to 93



**Diagnostic A205\_a (mission en)**


**Name :** Differences between maps of SLA (2)

**Input data :** Along track SLA


**Description :** The difference of SLA maps (mean, standard deviation, slope) is calculated from maps derived from diagnostic A203 using successively both altimetric components in the SLA calculation over a given period. This can be done globally, or separating in ascending and descending passes (except for SLA Grids).

Diagnostic type : Global internal analyses

SLA with GPD amplitude – SLA with RWT amplitude : annual signal  
Mission en, cycles 9 to 93



SLA with GPD phase – SLA with RWT phase : annual signal  
Mission en, cycles 9 to 93



## Diagnostic A205\_b (mission en)


**Name :** Differences between maps of SLA (2)

**Input data :** Along track SLA


**Description :** The difference of SLA maps (mean, standard deviation, slope) is calculated from maps derived from diagnostic A203 using successively both altimetric components in the SLA calculation over a given period. This can be done globally, or separating in ascending and descending passes (except for SLA Grids).

Diagnostic type : Global internal analyses

SLA with GPD amplitude – SLA with RWT amplitude : semi-annual signal  
Mission en, cycles 9 to 93



SLA with GPD phase – SLA with RWT phase : semi-annual signal  
Mission en, cycles 9 to 93





### Diagnostic A206\_a (mission en)

**Name :** Periodogram derived from temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The periodogram derived from temporal evolution of SLA (global, northern or southern hemisphere) can be done over all periods or focusing on particular periods, such as annual, semi annual or 60 day signal. Therefore mean of SLA differences are computed (every day or cycle), and time data series are plotted as a periodogram.

Diagnostic type : Global internal analyses





## Diagnostic A206\_b (mission en)

**Name :** Periodogram derived from temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA

**Description :** The periodogram derived from temporal evolution of SLA (global, northern or southern hemisphere) can be done over all periods or focusing on particular periods, such as annual, semi annual or 60 day signal. Therefore mean of SLA differences are computed (every day or cycle), and time data series are plotted as a periodogram.

Diagnostic type : Global internal analyses



## Diagnostic A206\_c (mission en)


**Name :** Periodogram derived from temporal evolution of Sea Level Anomaly (SLA)

**Input data :** Along track SLA


**Description :** The periodogram derived from temporal evolution of SLA (global, northern or southern hemisphere) can be done over all periods or focusing on particular periods, such as annual, semi annual or 60 day signal. Therefore mean of SLA differences are computed (every day or cycle), and time data series are plotted as a periodogram.

Diagnostic type : Global internal analyses

Periodogram of south hemisphere SLA (reference period = 1 year)  
Mission en, cycles 9 to 93



Periodogram of south hemisphere SLA (period = [0, 1 year])  
Mission en, cycles 9 to 93





## Diagnostic A207 (mission en)

**Name :** Sea Level Anomaly (SLA) versus coastal distance

**Input data :** Along track SLA

**Description :** Mean and standard deviation of SLA - computed by using successively both altimetric components - are plotted in function of coastal distances between 0 and 100 km.

Diagnostic type : Global internal analyses



**Diagnostic A208 (mission en)**


**Name :** Sea Level Anomaly (SLA) differences versus coastal distance

**Input data :** Along track SLA

**Description :** The differences of SLA variances - computed by using successively both altimetric components - are plotted in function of coastal distances between 0 and 100 km.

Diagnostic type : Global internal analyses

VAR(SLA with GPD) - VAR(SLA with RWT)  
Mission en, cycles 9 to 93



**Diagnostic A209 (mission en)**


**Name :** Differences between maps of SLA (3)

**Input data :** Along track SLA

**Description :** The differences between maps of SLA are calculated from the SLA differences (mean, standard deviation) using successively both altimetric components in the SLA calculation.

Diagnostic type : Global internal analyses

**VAR(SLA with GPD) – VAR(SLA with RWT)**  
**Mission en, cycles 9 to 93**



### Diagnostic C001 (mission en)


**Name :** Temporal evolution of SSH differences between tide gauges and altimetry measurements

**Input data :** Tide gauges SSH measurements

**Description :** The temporal evolution of global statistics (mean, variance, slope) of SSH differences between tide gauges and altimeter measurements are calculated from a cyclic way (altimeter repetivity) using successively both altimetric components in SSH calculation. The altimetric and tide gauges data are collocated with criteria of maximum of correlation, and tide gauges used are derived from global networks (GLOSS/CLIVAR, REFMAR).

Diagnostic type : Altimetry and in-situ data comparison

SLA differences : altimetry measurements - tide gauges  
Mission en, cycles 9 to 93



### Diagnostic C002 (mission en)


**Name :** Differences of temporal evolution of SSH differences between tide gauges and altimetry measurements

**Input data :** Tide gauges SSH measurements

**Description :** The difference between temporal evolution of global statistics of differences between tide gauge and altimeter data differences are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in altimetric SSH calculation. The altimetric and tide gauges data are collocated with criteria of maximum of correlation, and tide gauges used are derived from global networks as GLOSS/CLIVAR.

Diagnostic type : Altimetry and in-situ data comparison

Difference of variances :  $\text{VAR}(\text{SLA with GPD} - \text{T. G.}) - \text{VAR}(\text{SLA with RWT} - \text{T. G.})$   
Mission en, cycles 9 to 93




### Diagnostic C003 (mission en)

**Name :** Periodogram derived from temporal evolution of SSH differences between tide gauges and altimetry


**Input data :** Tide gauges SSH measurements

**Description :** The periodogram derived from temporal evolution of altimetric and tide gauges SSH differences is calculated using successively both altimetric components in the altimetric SSH. The periodogram is calculated from the mean or variance statistics and it can be displayed for all the whole time period or a dedicated one

Periodogram of SLA differences : altimetry measurements - tide gauges (ref. period = 1 year)  
Mission en, cycles 9 to 93



Periodogram of SLA differences : altimetry measurements - tide gauges (period = [0, 1 year])  
Mission en, cycles 9 to 93



**Diagnostic C004 (mission en)**

**Name :** Histograms of differences between tide gauges and altimeter SSH differences

**Input data :** Tide gauges SSH measurements

**Description :** The difference of histograms between altimeter and tide gauge SSH differences is computed from the elementary statistics (mean, variance) at each tide gauge using successively both altimetric components in the altimetry SSH.

Diagnostic type : Altimetry and in-situ data comparison

Histogram of the difference of variances :  $\text{VAR}(\text{SLA with GPD} - \text{T. G.}) - \text{VAR}(\text{SLA with RWT} - \text{T. G.})$   
Mission en, cycles 9 to 93

