Validation of new Ku band ionospheric instrumental correction after reversing the S-band PTR waveform

Study variable	**IONO_FILTR_ITERCorr2**
Reference variable	**IONO_FILTR_ITER**
Missions	Envisat *(en)*
Period	[19126, 21224]

Creation date : 2015/07/06

Contents

A000 - Differences of number of valid and invalid measurements between both altimetric components 3

A001 - Maps of differences of valid and invalid measurements between both altimetric components 4

A002 - Map of differences between both altimetric components over all the period 6

A003 - Periodogram derived from temporal evolution of altimetric component differences 7

A004 - Altimetric component differences versus coastal distances, latitude and longitude 9

A101 - Temporal evolution of SSH crossovers 12

A102 - Differences between temporal evolution of SSH crossovers 14

A103 - Map of SSH crossovers 15

A104 - Differences between maps of SSH crossovers 16

A105 - Differences between SSH crossovers vs coastal distance 17

A201 - Temporal evolution of Sea Level Anomaly (SLA) 18

A202 - Differences between temporal evolution of Sea Level Anomaly (SLA) 24

A203 - Map of Sea Level Anomaly (SLA) over all the period 26

A204 - Differences between maps of SLA trends 29

A205 - Differences between maps of SLA amplitude and phase 31

A206 - Periodogram derived from temporal evolution of Sea Level Anomaly (SLA) 33

A207 - Sea Level Anomaly (SLA) versus coastal distance 36

A208 - Sea Level Anomaly (SLA) differences versus coastal distance, latitude and longitude 37

A209 - Differences between maps of SLA variance 39

A210 - Differences between maps of SLA variance for different frequency bands 40

A211 - Differences between maps of SLA per year 43

Diagnostic type : Mono-mission analyses

Diagnostic A000 (mission en)

Name : Differences of number of valid and invalid measurements between both altimetric components

Input data : Along track altimetric components

Description : The number of valid measurements for one parameter and invalid for the other, and vice-versa.

Diagnostic type : Mono-mission analyses

Diagnostic A001 (mission en)

Name : Maps of differences of valid and invalid measurements between both altimetric components

Input data : Along track altimetric components

Description : The first map represents the valid measurements for one parameter and invalid for the other, and vice-versa for the second map.

Diagnostic type : Mono-mission analyses

Diagnostic A001 (mission en)

Name : Temporal evolution of differences between both altimetric components

Input data : Along track altimetric components

Description : The temporal evolution of global statistics (mean, variance, slope) of differences between 2 different standards of a same altimetric component (sea surface height correction, altimeter parameter, orbit) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) . These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Mono-mission analyses

Diagnostic A002 (mission en)

Name : Map of differences between both altimetric components over all the period

Input data : Along track altimetric components

Description : The map of global statistics (mean, standard deviation) of differences between 2 different standards of a same altimetric component (sea surface height correction, altimeter parameter, orbit) are calculated over a given period which is the longer as possible to have obtain reliable statically results. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Mono-mission analyses

Diagnostic A003_a (mission en)

Name : Periodogram derived from temporal evolution of altimetric component differences

Input data : Along track altimetric components

Description : The periodogram derived from temporal and global altimetric component differences is calculated from cycle by cycle monitoring of altimetric component differences (derived from diagnostic A001). It is calculated from the mean or the variance differences. The Periodogram can be calculated for all the periods, but it can be focused on a dedicated period.

Diagnostic type : Mono-mission analyses

Diagnostic A003_b (mission en)

Name : Periodogram derived from temporal evolution of altimetric component differences

Input data : Along track altimetric components

Description : The periodogram derived from temporal and global altimetric component differences is calculated from cycle by cycle monitoring of altimetric component differences (derived from diagnostic A001). It is calculated from the mean or the variance differences. The Periodogram can be calculated for all the periods, but it can be focused on a dedicated period.

Diagnostic type : Mono-mission analyses

Diagnostic A004 (mission en)

Name : Altimetric component differences versus coastal distances, latitude and longitude

Input data : Along track altimetric components

Description : Mean and standard deviation of the differences between 2 different standards of a same altimetric component (sea surface height correction, altimeter parameter, orbit) are computed and plotted in function of coastal distances between 0 and 100 km, in function of latitudes and in fonction of longitudes.

Diagnostic type : Mono-mission analyses

Diagnostic A004 (mission en)

Name : Altimetric component differences versus coastal distances, latitude and longitude

Input data : Along track altimetric components

Description : Mean and standard deviation of the differences between 2 different standards of a same altimetric component (sea surface height correction, altimeter parameter, orbit) are computed and plotted in function of coastal distances between 0 and 100 km, in function of latitudes and in fonction of longitudes.

Diagnostic type : Mono-mission analyses

Diagnostic A004 (mission en)

Name : Altimetric component differences versus coastal distances, latitude and longitude

Input data : Along track altimetric components

Description : Mean and standard deviation of the differences between 2 different standards of a same altimetric component (sea surface height correction, altimeter parameter, orbit) are computed and plotted in function of coastal distances between 0 and 100 km, in function of latitudes and in fonction of longitudes.

Diagnostic type : Mono-mission analyses

Diagnostic A101_a (mission en)

Name : Temporal evolution of SSH crossovers

Input data : Sea Surface Height (SSH) crossovers

Description : The temporal evolution of global statistics (mean, standard deviation) of SSH differences are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Mono-mission analyses

Diagnostic A101_b (mission en)

Name : Temporal evolution of SSH crossovers

Input data : Sea Surface Height (SSH) crossovers

Description : The temporal evolution of global statistics (mean, standard deviation) of SSH differences are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Mono-mission analyses

Diagnostic A102 (mission en)

Name : Differences between temporal evolution of SSH crossovers

Input data : Sea Surface Height (SSH) crossovers

Description : The difference of temporal evolution between the global statistics (mean, standard deviation) of SSH differences are calculated using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Mono-mission analyses

Diagnostic A103 (mission en)

Name : Map of SSH crossovers

Input data : Sea Surface Height (SSH) crossovers

Description : The differences between maps of SSH crossovers differences (mean, variance) are calculated using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Mono-mission analyses

Diagnostic A104 (mission en)

Name : Differences between maps of SSH crossovers

Input data : Sea Surface Height (SSH) crossovers

Description : The differences between maps of SSH crossovers (derived from diagnostic A103) are calculated from the SSH crossover differences (mean, standard deviation) using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Mono-mission analyses

Diagnostic A105 (mission en)

Name : Differences between SSH crossovers vs coastal distance

Input data : Sea Surface Height (SSH) crossovers

Description : The differences of SSH variances at crossovers are plotted in function of coastal distance, latitudes and longitudes.

Diagnostic type : Mono-mission analyses

Diagnostic A201_a (mission en)

Name : Temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes, or separating North and South hemispheres.

Diagnostic type : Mono-mission analyses

Diagnostic A201_b (mission en)

Name : Temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes, or separating North and South hemispheres.

Diagnostic type : Mono-mission analyses

Diagnostic A201_c (mission en)

Name : Temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes, or separating North and South hemispheres.

Diagnostic type : Mono-mission analyses

Diagnostic A201_d (mission en)

Name : Temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes, or separating North and South hemispheres.

Diagnostic type : Mono-mission analyses

Diagnostic A201_e (mission en)

Name : Temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes, or separating North and South hemispheres.

Diagnostic type : Mono-mission analyses

Diagnostic A201_f (mission en)

Name : Temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes, or separating North and South hemispheres.

Diagnostic type : Mono-mission analyses

Diagnostic A202_a (mission en)

Name : Differences between temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The differences between temporal evolution of SLA are calculated from statistics derived from diagnostic A201 (mean, variance) using 2 different components in the SLA calculation. They are calculated globally, but also separating ascending and descending passes or separating North and South hemispheres.

Diagnostic type : Mono-mission analyses

Diagnostic A202_b (mission en)

Name : Differences between temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The differences between temporal evolution of SLA are calculated from statistics derived from diagnostic A201 (mean, variance) using 2 different components in the SLA calculation. They are calculated globally, but also separating ascending and descending passes or separating North and South hemispheres.

Diagnostic type : Mono-mission analyses

Diagnostic A203_a (mission en)

Name : Map of Sea Level Anomaly (SLA) over all the period

Input data : Along track SLA

Description : The map of global statistics (mean, standard deviation) of SLA are calculated using successively both altimetric components in the SLA calculation over a large period. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Mono-mission analyses

Diagnostic A203_b (mission en)

Name : Map of Sea Level Anomaly (SLA) over all the period

Input data : Along track SLA

Description : The map of global statistics (mean, standard deviation) of SLA are calculated using successively both altimetric components in the SLA calculation over a large period. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Mono-mission analyses

Diagnostic A203_c (mission en)

Name : Map of Sea Level Anomaly (SLA) over all the period

Input data : Along track SLA

Description : The map of global statistics (mean, standard deviation) of SLA are calculated using successively both altimetric components in the SLA calculation over a large period. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Mono-mission analyses

Diagnostic A204_a (mission en)

Name : Differences between maps of SLA trends

Input data : Along track SLA

Description : The difference of SLA maps (mean, standard deviation, slope) is calculated from maps derived from diagnostic A203 using successively both altimetric components in the SLA calculation over a given period. This can be done globally, or separating in ascending and descending passes (except for SLA Grids).

Diagnostic type : Mono-mission analyses

Diagnostic A204_b (mission en)

Name : Differences between maps of SLA trends

Input data : Along track SLA

Description : The difference of SLA maps (mean, standard deviation, slope) is calculated from maps derived from diagnostic A203 using successively both altimetric components in the SLA calculation over a given period. This can be done globally, or separating in ascending and descending passes (except for SLA Grids).

Diagnostic type : Mono-mission analyses

Diagnostic A205_a (mission en)

Name : Differences between maps of SLA amplitude and phase

Input data : Along track SLA

Description : The difference of SLA maps (mean, standard deviation, slope) is calculated from maps derived from diagnostic A203 using successively both altimetric components in the SLA calculation over a given period. This can be done globally, or separating in ascending and descending passes (except for SLA Grids).

Diagnostic type : Mono-mission analyses

Diagnostic A205_b (mission en)

Name : Differences between maps of SLA amplitude and phase

Input data : Along track SLA

Description : The difference of SLA maps (mean, standard deviation, slope) is calculated from maps derived from diagnostic A203 using successively both altimetric components in the SLA calculation over a given period. This can be done globally, or separating in ascending and descending passes (except for SLA Grids).

Diagnostic type : Mono-mission analyses

Diagnostic A206_a (mission en)

Name : Periodogram derived from temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The periodogram derived from temporal evolution of SLA (global, northern or southern hemisphere) can be done over all periods or focusing on particular periods, such as annual, semi annual or 60 day signal. Therefore mean of SLA differences are computed (every day or cycle), and time data series are plotted as a periodogram.

Diagnostic type : Mono-mission analyses

Diagnostic A206_b (mission en)

Name : Periodogram derived from temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The periodogram derived from temporal evolution of SLA (global, northern or southern hemisphere) can be done over all periods or focusing on particular periods, such as annual, semi annual or 60 day signal. Therefore mean of SLA differences are computed (every day or cycle), and time data series are plotted as a periodogram.

Diagnostic type : Mono-mission analyses

Diagnostic A206_c (mission en)

Name : Periodogram derived from temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The periodogram derived from temporal evolution of SLA (global, northern or southern hemisphere) can be done over all periods or focusing on particular periods, such as annual, semi annual or 60 day signal. Therefore mean of SLA differences are computed (every day or cycle), and time data series are plotted as a periodogram.

Diagnostic type : Mono-mission analyses

Diagnostic A207 (mission en)

Name : Sea Level Anomaly (SLA) versus coastal distance

Input data : Along track SLA

Description : Mean and standard deviation of SLA - computed by using successively both altimetric components - are plotted in function of coastal distances between 0 and 100 km.

Diagnostic type : Mono-mission analyses

Diagnostic A208 (mission en)

Name : Sea Level Anomaly (SLA) differences versus coastal distance, latitude and longitude

Input data : Along track SLA

Description : The differences of SLA variances - computed by using successively both altimetric components - are plotted in function of coastal distances between 0 and 100 km, in function of latitudes and in function of longitudes.

Diagnostic type : Mono-mission analyses

Diagnostic A208 (mission en)

Name : Sea Level Anomaly (SLA) differences versus coastal distance, latitude and longitude

Input data : Along track SLA

Description : The differences of SLA variances - computed by using successively both altimetric components - are plotted in function of coastal distances between 0 and 100 km, in function of latitudes and in function of longitudes.

Diagnostic type : Mono-mission analyses

Diagnostic A209 (mission en)

Name : Differences between maps of SLA variance

Input data : Along track SLA

Description : The differences between maps of SLA are calculated from the SLA differences (mean, standard deviation) using successively both altimetric components in the SLA calculation.

Diagnostic type : Mono-mission analyses

Diagnostic A210_a (mission en)

Name : Differences between maps of SLA variance for different frequency bands

Input data : Along track SLA

Description : The differences between maps of SLA (variance) are calculated from the mean SLA maps using successively both altimetric components in the SLA calculation filtered to separate high-frequency (T < 1 yr), mid-frequency (1 yr < T < 3 yrs) and low-frequency (T > 3 yrs) signals.

Diagnostic type : Mono-mission analyses

Diagnostic A210_b (mission en)

Name : Differences between maps of SLA variance for different frequency bands

Input data : Along track SLA

Description : The differences between maps of SLA (variance) are calculated from the mean SLA maps using successively both altimetric components in the SLA calculation filtered to separate high-frequency (T < 1 yr), mid-frequency (1 yr < T < 3 yrs) and low-frequency (T > 3 yrs) signals.

Diagnostic type : Mono-mission analyses

Diagnostic A210_c (mission en)

Name : Differences between maps of SLA variance for different frequency bands

Input data : Along track SLA

Description : The differences between maps of SLA (variance) are calculated from the mean SLA maps using successively both altimetric components in the SLA calculation filtered to separate high-frequency (T < 1 yr), mid-frequency (1 yr < T < 3 yrs) and low-frequency (T > 3 yrs) signals.

Diagnostic type : Mono-mission analyses

Diagnostic A211 (mission en)

Name : Differences between maps of SLA per year

Input data : Along track SLA

Description : The differences between map of SLA (mean) are calculated for each year using successively both altimetric components in the SLA calculation